

conditions / S. Vojinovic et al. *Vojnosanitetski pregled*. 2014. Vol. 72. P. 30-30.

DOI: <https://doi.org/10.2298/VSP140121030V>

11. Dunn S. E., Lee H., Pavri F. R., Zhang M. A. Sex-Based Differences in Multiple Sclerosis (Part I): Biology of Disease Incidence. *Current Topics in Behavioral Neurosciences*. Springer. 2015. Vol. 26. P. 29-56. DOI: https://doi.org/10.1007/7854_2015_371

12. Gomez F. P., Steelman A. J., Young C. R., Welsh C. J. Hormone and immune system interactions in demyelinating disease. *Hormones and Behavior*. 2013. Vol. 63, No. 2. P. 315-321.

DOI: <https://doi.org/10.1016/j.yhbeh.2012.10.014>

13. Seasonal variations of 25-OH vitamin D serum levels are associated with clinical disease activity in multiple sclerosis patients / C. Hartl et al. *Journal of the Neurological Sciences*. 2017. Vol. 375. P. 160-164.

DOI: <https://doi.org/10.1016/j.jns.2017.01.059>

14. Sex hormones modulate brain damage in multiple sclerosis: MRI evidence / V. Tomassini et al. *J. neurology, neurosurgery, and psychiatry*. 2005. Vol. 76. P. 272-275. DOI: <https://doi.org/10.1136/jnnp.2003.033324>

15. Sicotte N. L., Giesser B. S., Tandon V. Testosterone treatment in multiple sclerosis: A pilot study. *Arch Neurol*. 2007. Vol. 64. P. 683-688.

DOI: <https://doi.org/10.1001/archneur.64.5.683>

Стаття надійшла до редакції
20.06.2019



UDC 616.972-036.15-085-097

<https://doi.org/10.26641/2307-0404.2019.3.181889>

S.V. Zakharov,
V.K. Zakharov,
V.V. Gorbuntsov

THE CONTENT OF CYTOKINES IN THE BLOOD SERUM OF PATIENTS WITH EARLY LATENT SYPHILIS IN PROCESS OF TREATMENT

SE «Dnipropetrovsk medical academy of Health Ministry of Ukraine»

Department of skin and venereal diseases

V. Vernadsky str., 9, Dnipro, 49044, Ukraine

ДЗ «Дніпропетровська медична академія МОЗ України»

кафедра шкірних та венеричних хвороб

(зав. – д. мед. н., проф. А.Д. Дюдюн)

вул. В. Вернадського, 9, Дніпро, 49044, Україна

e-mail: dsma@dsma.dp.ua

Цитування: *Медичні перспективи*. 2019. Т. 24, № 3. С. 96-101

Cited: *Medicni perspektivi*. 2019;24(3):96-101

Key words: *syphilis latent early, cytokines, treatment, prognosis, serological resistance*

Ключові слова: *сифіліс прихований ранній, цитокіни, лікування, прогноз*

Ключевые слова: *сифилис скрытый ранний, цитокины, лечение, прогноз*

Abstract. **The content of cytokines in the blood serum of patients with early latent syphilis in process of treatment.** Zakharov S.V., Zakharov V.K., Gorbuntsov V.V. *Objective – to study the concentration of pro- and anti-inflammatory cytokines in patients with early latent syphilis before and after treatment. The study was conducted in 112 patients with early latent syphilis (52 men and 60 women) and 15 healthy persons of the control group – all aged 18-43. Serological tests were used: the classical complex of serological reactions (CSR), ELISA, the reaction of passive hemagglutination (RPGA), immunofluorescence reaction with absorption (RIF-abs) and RIF-200. Using ELISA, the levels of cytokines IL-2, IL-6, IL-10, TNF α , and INF γ were determined in serum (in patients with latent syphilis, before and after treatment). The analysis of the received data was carried out with application of the program package Statistics 6.0. To identify the relationships between the indices, Friedman's nonparametric variance analysis with the*

definition of χ^2 was used. The concentration of IL-10 before treatment in patients with early latent syphilis was significantly increased by 3.7 (14.9 ± 0.9 pg/ml compared to the control group 4.11 ± 0.5 pg/ml). The dependence of the content of this cytokine on the period of infection was established. It was found that one year after treatment concentration of IL-10 remained 1.8 times higher. The concentration of IL-6 was increased by 6.5 times in patients with latent early syphilis and was also dependent on the period of the infection, and also even 1.5-2 years after treatment the IL-6 content remained elevated. It was also found that in patients with latent early syphilis the concentration of IL-2 was increased by 2.3 times and it did not normalize in 24.5% of patients even two years after treatment. In patients with latent early syphilis, the concentration of TNF α was increased by 3.6 times. After treatment, the concentration of TNF α decreased, but exceeded the control values by 1.55 times. Further analysis of the concentration of TNF α revealed that, with a disease period of up to one year, the concentration of this cytokine was normal during ten months after treatment, and in patients with the disease period of more than one year, after treatment normalization of the TNF α level was absent in 64.5% even after 18 months. It was found that the concentration of INF γ in patients with early latent syphilis before treatment was increased by 3.8 times in comparison with the control group. After treatment, almost 25 % of patients with the duration of the infection more than one year revealed no normalization of INF γ and it was increased by 1.5 times. In patients with latent early syphilis there was a significant increase in the concentration of cytokines TNF α and INF γ , an imbalance of IL-2, IL-6, IL-10 as well as. The dependence of the cytokine concentration on the period of infection is noted. Based on the study of the cytokine status in patients with early latent syphilis with periods of infection of more than one year, the use of immunomodulatory therapy may be recommended. Increasing the concentration of such cytokines as TNF α , IL-6, IL-10 after treatment can be used as prognostic tests of serological resistance.

Реферат. Вміст цитокінів у сироватці крові хворих на ранній прихований сифіліс у процесі лікування. Захаров С.В., Захаров В.К., Горбунцов В.В. Мета роботи - вивчити концентрацію про- і протизапальних цитокінів у пацієнтів з раннім прихованим сифілісом до і після лікування. Дослідження проводилося у 112 пацієнтів з раннім прихованим сифілісом (жінок - 60, чоловіків - 52) і 15 здорових пацієнтів групи контролю віком 18-43 роки. Застосовували серологічні методи дослідження: РЗК, ІФА, РПГА, РІФ. Методом ІФА визначали в сироватці крові (у пацієнтів з прихованим сифілісом - до і після лікування) рівні цитокінів IL-2, IL-6, IL-10, TNF α , і INF γ . Аналіз отриманих даних проводився із застосуванням пакета програм Статистика 6.0. Для виявлення зв'язків між показниками застосовували непараметричний дисперсійний аналіз Фрідмана з визначенням χ^2 . Концентрація IL-10 до лікування в пацієнтів з раннім прихованим сифілісом була достовірно підвищена в 3,7 раза ($14,9 \pm 0,9$ пг/мл порівняно з групою контролю $4,11 \pm 0,5$ пг/мл). Було встановлено залежність вмісту цього цитокіну від термінів інфекції. Було встановлено, що через один рік після лікування концентрація IL-10 залишалася підвищеною в 1,8 раза. Концентрація IL-6 була підвищена в пацієнтів з прихованим раннім сифілісом у 6,5 раза і також була залежна від термінів інфекції, і навіть через 1,5-2 роки після лікування вміст IL-6 залишався підвищеним. Було також встановлено, що в пацієнтів з прихованим раннім сифілісом концентрація IL-2 була підвищена в 2,3 раза і вона не нормалізувалася в 24,5% пацієнтів навіть через два роки після лікування. У пацієнтів з прихованим раннім сифілісом концентрація TNF α була підвищена в 3,6 раза. Після проведеного лікування концентрація TNF α у них зменшувалася, але перевищувала показники контролю в 1,55 раза. При подальшому аналізі концентрації TNF α було встановлено, що при термінах захворювання до одного року через десять місяців після лікування була нормалізація концентрації цього цитокіну, а в пацієнтів з термінами захворювання більше одного року після лікування нормалізації рівня TNF α не було в 64,5% навіть через 18 місяців. Було встановлено, що концентрація INF γ у пацієнтів з раннім прихованим сифілісом до лікування була підвищена в 3,8 раза порівняно з контрольною групою. Після лікування майже в 25% пацієнтів з термінами інфекції більше одного року нормалізації INF γ не було і він був підвищеним в 1,5 рази. У пацієнтів з прихованим раннім сифілісом спостерігається значне підвищення концентрації цитокінів TNF α і INF γ , а також дисбаланс IL-2, IL-6, IL-10. Відзначається залежність концентрації цитокінів від термінів інфікування. На підставі вивчення цитокінового статусу в пацієнтів з раннім прихованим сифілісом з термінами інфікування більше одного року можна рекомендувати застосування імуномодуючої терапії. Підвищення концентрації таких цитокінів, як TNF α , IL-6, IL-10, після лікування може бути використано в якості прогностичних тестів серологічної резистентності.

The urgency of the latent syphilis problem is caused by the fact that in the structure of syphilis incidence its part amounts more than 50 %, the mechanisms of its pathogenesis remain unstudied completely, and after the treatment of such patients the serological resistance arises [1, 3, 4, 7, 9].

Among numerous factors contributing to the increase of the number of cases of the early latent syphilis, a significant part is belongs to [1, 2, 4, 11]:

- the significant spread of antibiotics using, both for the treatment of syphilis and for the treatment of other STIs;
- expansion and introduction into practice of a number of highly sensitive methods of examination (immunoenzyme method, immune blotting, the reaction of passive hemagglutination (RPGA));
- the reduction of the virulent properties of the pale treponema itself and its resistance to therapy.

In addition, some researchers believe that *L*- and cyst-forms of the pale treponema itself can be one of the reasons of the latent course of syphilis [4, 18]. At the same time, none of the researchers studying the mechanisms of the syphilitic infection pathogenesis has doubts as to the fact that the prognosis of the disease depends on the state of the immune system [7, 12, 15, 16, 17]. Until now, there is no consensus among specialists on understanding the mechanism of immunity disorders under the latent syphilis [2, 4, 5].

The above data indicate that the issue of the classification and mechanisms of the immune system disorders initiation under this form of syphilis remains unresolved [2, 16, 17]. Studies of the content of anti-inflammatory cytokines in patients with early latent syphilis, which have been conducted earlier, are highly contradictory and incomplete [4, 5, 8-13].

Under the chronic inflammation peculiar to syphilitic infection in particular, the normal cytokine profile changes, and that is manifested by increased secretion of pro-inflammatory cytokines [3, 5, 6].

However there is another view on the role of interleukins created in the cells of a sick person under the influence of pale treponema due to the activation of *Th1* cells [12, 13].

To this day almost no studies have been conducted on the content of *IL-2* in the blood serum of patients with latent syphilis, while *IL-2* plays a significant role in the activation of the immune system through the stimulation of the synthesis of *INF γ* , *IL-6*. Unlike *IL-2*, *IL-6* is produced by different types of cells by activation of *CD4+* and *CD8+* and belongs to the proinflammatory cytokines. One of the most important cytokines determining the immune response is *IL-10*. Under its influence, cellular immunity (*Th1*) is inhibited and humoral immunity (*Th2*) is stimulated, the secretion of *INF γ* , *TNF α* , *IL-1*, *IL-8* decreases. *TNF α* is a polypotent immunomodulatory cytokine, namely a mediator of a specific and non-specific immune response that depends on its concentration. *TNF α* is capable of activating *EK*-cells, increasing the secretion of *IL-6*. *TNF α* is synthesized mainly by *Th1*-lymphocytes and *EK*-cells and has a significant antibacterial and immunomodulatory effect [5, 6].

A comprehensive study of the cytokine profile in patients with early latent syphilis before and after treatment can give a more detailed idea of the mechanism of the immune response and allow further prognosis of the disease [10, 19-22].

Objective – to study the concentration of pro- and anti-inflammatory cytokines in patients with latent early syphilis before and after treatment.

MATERIALS AND METHODS OF RESEARCH

The study was conducted in 112 patients (52 men and 60 women) aged 18-43 (mean age 31.1±0.7 years) with early latent syphilis who were on treatment at the skin and venereal disease clinic of the State Establishment "Dnipropetrovsk Medical Academy of Health Ministry of Ukraine". The comparison group consisted of 15 practically healthy persons with similar sex and age distribution.

The criteria for inclusion in the study were: age up to 45 years, diagnosis of early latent syphilis. The criteria for exclusion from the study were: age over 45, pregnancy, presence of tuberculosis, viral hepatitis, HIV / AIDS, other infectious diseases; diabetes, as well as antibiotic treatment in the last three months before the study.

For the diagnosis of syphilis serological tests were used: the classical complex of serological reactions (CSR), ELISA (*IgM*, *IgG*), the reaction of passive hemagglutination (RPGA), immunofluorescence reaction with absorption (RIF-abs) and RIF-200. The serum levels of *IL-2*-, *IL-6*-, *IL-10*-cytokone, *TNF α* , and *INF γ* were measured by the method of enzyme immunoassay [11]. The analysis of the data received was carried out with application of the program package Statistics 6.0. The difference in indices with $p < 0.05$ considered as probable one. To identify the relationships between the indices, Friedman's nonparametric variance analysis with the definition of χ^2 was used. The coupling between indices was considered significant if χ^2 value exceeded the critical one ($\chi^2 = 3.84$) [3, 4].

RESULTS AND DISCUSSION

The concentration of *IL-10* in the blood serum of patients with latent early syphilis before treatment was probably ($p < 0.05$) increased to 14.9±0.9 pg/ml in comparison with the control group (4.11±0.5 pg/ml), that is by 3.7 times. When analyzing this anti-inflammatory interleukin in patients with an infection period of more than one year, its concentration was 5.7 times higher than in healthy persons (23.5±0.6 and 4.11±0.5; $p < 0.05$). If before treatment the concentration of *IL-10* was increased by 3.7 times, after treatment the content of this interleukin significantly decreased to 10.1±0.2 pg/ml and exceeded the normal values by 2.4 ($p < 0.05$). On further observation after treatment, the further decrease in *IL-10* concentration to 7.5±0.1 pg/ml was marked only after 12.3±1.2 months. That is, even one year after treatment there was no normalization of the *IL-10* concentration (it remained 1.8 times higher than one in the control group).

In patients with early latent syphilis before treatment, the concentration of *IL-6* was 31.3±1.2 pg/ml,

that is it was increased by 6.5 times in comparison with the control group.

When analyzing the concentration of *IL-6*, it was found that in patients with a period of infection up to one year it exceeded the norm by almost 10 times (48.3 ± 1.5 compared to 4.8 ± 0.2 , $p < 0.05$). Under the longer duration of the disease, the concentration of *IL-6* was almost three times less (15.6 ± 0.5 pg/ml).

After treatment the serum concentration of *IL-6* remained elevated (7.4 ± 0.8 pg/ml, $p < 0.05$) even 1.5-2.0 years after.

The concentration of *IL-2* in the blood serum of patients with latent early syphilis before treatment was 37.8 ± 4.1 pg/ml while a control group score was 16.5 ± 3.2 pg/ml, that is, it was increased by 2.3 ($p < 0.05$) times. When analyzing the concentration of this cytokine depending on the duration of the disease to one year and more than one year, the difference was unreliable. After treatment the normalization of *IL-2* secretion was lacking in 24.5% of patients even two years after.

When analyzing the secretion of tumor necrosis factor (TNF α), we obtained such results: in patients with latent early syphilis the concentration of TNF α was 18.8 ± 0.6 pg/ml (in the control group 5.2 ± 0.5 pg/ml, $p < 0.05$), that is, it was increased by 3.6 times. After treatment, the concentration of TNF α decreased to 8.1 ± 0.3 pg/ml ($p < 0.05$) and exceeded the control values by 1.55 times.

Further analysis of the concentration of TNF α revealed that, under the disease period of up to one year, the concentration of this cytokine was normal

during ten months after treatment, and in patients with a disease period of more than one year normalization of the TNF α level after treatment was lacking in 64.5% even 18 months after.

The concentration of INF γ in patients with early latent syphilis before treatment was increased by 3.8 times in comparison with the control group (53.7 ± 3.1 pg/ml vs. 16.1 ± 1.3 , $p < 0.05$). After treatment almost 75% of patients revealed normalization of this cytokine and only 25% of patients with period of infection more than one year, revealed no normalization of INF γ and it was increased by 1.5 times.

CONCLUSIONS

1. Under the early latent syphilis there is a significant increase in the concentration of cytokines that are synthesized Th1 type of the (TNF α , INF γ), as well as the imbalance of *IL-2*, *IL-6*, *IL-10*.

2. The dependence of the cytokines concentration on the terms of infection was established; so under duration of the infection of more than one year the imbalance of pro- and anti-inflammatory cytokines was more pronounced.

3. Based on the study of the cytokine status, patients with early latent syphilis with terms of infection of more than one year should be treated with medications causing a positive effect on the immune system.

4. An increase in the concentration of such cytokines as TNF α , *IL-6*, *IL-10* after treatment can be used as prognostic tests of serological resistance.

REFERENCES

- Bondarenko HM, Unuchko SV, Hubenko TV, Matyushenko VP. [Clinical and epidemiological features of syphilis at the present stage]. *Dermatolohiia ta venerolohiia*. 2014;2(64):65-71. Ukrainian.
- Bondarenko GM, Unuchko SV, Nikitenko IN, Gubenko TV, Kutovaia VV. [Syphilis: the current state of the problem]. *Dermatolohiia ta venerolohiia*. 2018;1(79):8-12. Russian.
- Burmeister GR, Pecutto A, Ulrichs T, Aikher A. [Visual immunology]. Moskva, Binom. 2014;320. Russian.
- Dmitriev GA, Dolya OV. [Diagnosis of syphilitic infection]. *Klinicheskaia dermatologiya i venerologiya*. 2012;6:10-15. Russian.
- Drannik GN. [European Declaration of Immunotherapy]. *Klinichna imunolohiia. Alerholohiia. Infektolohiia*. 2011;6-7:5-13. Russian.
- Drannik GN. [Modern ideas about the mechanisms of innate and acquired immunity and their interaction (part 1)]. *Liky Ukrainy*. 2013;4:22-29. Russian.
- Zakharov VK, Dyudyun AD, Zakharov SV. [Hidden syphilis]. *Dnipropetrovsk*. 2011;119. Ukrainian.
- Zakharov SV, Zakharov VK, Djudjun AD. [Features of the level of interleukin-6 in patients with latent syphilis]. Conference "Dermatovenerology in the aspect of a family doctor". Kyiv. 2008;35. Ukrainian.
- Zakharov SV. [Early latent syphilis: focuses on social, epidemiological and medical factors]. *Dermatolohiia ta venerolohiia*. 2018;1(79):44-50. Ukrainian.
- Zakharov SV. [Cytokine profile in patients with early latent syphilis]. *Medicni perspektivi*. 2018;23(4):71-5. Ukrainian. doi: <https://doi.org/10.26641/2307-0404.2018.1.124933>
- Kamyshnikov VS. [Clinical laboratory diagnostics (methods and interpretation of laboratory studies)]. Moskva, MEDpres-inform. 2015;720. Russian.
- Mavrov GI. [Outpatient treatment of patients with early syphilis with durant penicillins in combination with doxycycline]. *Dermatovenerologiya. Kosmetologiya. Seksopatologiya*. 2016;1-4:214-9. Russian.
- Mavrov GI, Mironyuk VI, Shcherbakova JuV, Osinskaya TV. [Some indicators of cytokine status in

patients with syphilis of psychoactive substance users]. *Dermatolohiia ta venerolohiia*. 2018;4:13-16. Ukrainian.

14. Mintser OP, Voronenko JuV, Vlasov VV. [Processing of clinical and experimental data in medicine]. Kyiv, Vyshha shkola. 2003;350. Ukrainian.

15. Sivak VV, Chebotarev VV, Chebotareva NV. [Syphilis in penitentiary system institutions]. Stavropol. 2003;166. Russian.

16. Shcherbakova JuV. [Diagnosis, treatment and prevention of latent syphilis based on the study of epidemiology, serological and cytokine status.]. [dissertation] Kharkiv: Institute of Dermatology and Venereology. 2005;20. Ukrainian.

17. Shcherbakova JuV. [Features of the state of immunity in patients with latent syphilis]. *Dermatolohiia ta venerolohiia*. 2005;5(27):27-32. Russian.

18. Carlson JA, Dabiri G, Cribier B, Sell S. The immunopathobiology of syphilis: the manifestations and course of syphilis are determined by the level of delayed-

hype hypersensitivity. *Am. J. Dermatopathol*. 2011;33(5):433-60.

doi: <https://doi.org/10.1097/DAD.0b013e3181e8b587>

19. Dinarello C. Proinflammatory cytokines. *Chest*. 2000;118(9):503-8.

doi: <https://doi.org/10.1378/chest.118.2.503>

20. Kenyon C, Osbak KK, Crucitti T, Kestens L. The immunological response to syphilis differs by HIV status; a prospective observational cohort study. *BMC Infectious Diseases*. 2017;17:111-9.

doi: <https://doi.org/10.1186/s12879-017-2201-7>

21. Knudsen A, Benfield T, Kofoed K. Cytokine expression during syphilis infection in HIV-1-infected individuals. *Sex. Transm. Dis.* 2009;36(5):300-4. doi: <https://doi.org/10.1097/OLQ.0b013e318193ca26>

22. Lusiak M, Podwińska J. Interleukin 10 and its role in the regulation of the cell-mediated immune response in syphilis. *Arch. Immunol Ther. Exp.* 2001;49(6):417-21.

СПИСОК ЛІТЕРАТУРИ

1. Бондаренко Г. М., Унучко С. В., Губенко Т. В., Матюшенко В. П. Клініко-епідеміологічні особливості сифілісу на сучасному етапі. *Дерматологія та венерологія*. 2014. Т. 64, № 2. С. 65-71.

2. Бондаренко Г. М., Унучко С. В., Никитенко І. Н., Губенко Т. В. Сифіліс: сучасні проблеми. *Дерматологія та венерологія*. 2018. Т. 79, № 1. С. 8-12.

3. Бурмейстер Г.-Р., Пецутто А., Улрихс Т., Айхер А. Наглядная иммунология. Москва: Бинном, 2014. 320 с.

4. Дмитриев Г. А., Доля О. В. Диагностика сифилитической инфекции. *Клиническая дерматология и венерология*. 2012. № 6. С. 10-15.

5. Дранник Г. Н. Европейская декларация по иммунотерапии. *Клінічна імунологія. Алергологія. Інфектологія*. 2011. № 6-7. С. 5-13.

6. Дранник Г. Н. Современные представления о механизмах врожденного и приобретенного иммунитета и их взаимодействие (часть 1). *Лици України*. 2013. № 4. С. 22-29.

7. Захаров В. К., Дюдюк А. Д., Захаров С. В. Прихований сифіліс. Дніпропетровськ. 2011. 119 с.

8. Захаров С. В., Захаров В. К., Дюдюк А. Д. Особливості рівня інтерлейкіну-6 у хворих на прихований сифіліс. *Дерматовенерологія в аспекті сімейного лікаря: матеріали наук.-практ. конф.* Київ. 2008. С. 35.

9. Захаров С. В. Ранній прихований сифіліс: у фокусі соціальні, епідеміологічні та медичні чинники. *Дерматологія та венерологія*. 2018. Т. 79, № 1. С. 44-50.

10. Захаров С. В. Цитокиновий профіль у хворих на ранній прихований сифіліс. *Медичні перспективи*. 2018. Т. XXIII, № 4. С. 71-75. DOI: <https://doi.org/10.26641/2307-0404.2018.1.124933>

11. Камышников В. С. Клиническая лабораторная диагностика (методы и трактовка лабораторных исследований). Москва: МЕД-пресс-информ, 2015. 720 с.

12. Мавров Г. І. Амбулаторное лечение больных ранним сифилисом дюрантными пенициллинами в сочетании с доксициклином. *Дерматовенерология. Косметология. Сексопатология*. 2016. № 1-4. С. 214-219.

13. Мавров Г. І. Миронюк В. І., Щербакова Ю. В., Осінська Т. В. Окремі показники цитокинового статусу у хворих на сифіліс споживачів психоактивних речовин. *Дерматологія та венерологія*. 2018. № 4. С. 13-16.

14. Мінцер О. П., Вороненко Ю. В., Власов В. В. Оброблення клінічних і експериментальних даних у медицині: навч. посіб. Київ: Вища школа, 2003. 350 с.

15. Сивак В. В., Чеботарев В. В., Чеботарева Н. В. Сифіліс в учреждениях пенитенциарной системы. Ставрополь, 2003. 166 с.

16. Щербакова Ю. В. Диагностика, лікування та профілактика прихованого сифілісу на основі вивчення епідеміології, серологічного та цитокинового статусу: автореф. дис. ... канд. мед. наук: 14.01.20 / АМН України. Ін-т дерматології та венерології. Харків, 2005. 20 с.

17. Щербакова Ю. В. Особенности состояния иммунитета у больных скрытым сифилисом. *Дерматология та венерология*. 2005. Т. 27, № 5. С. 27-32.

18. Carlson J. A., Dabiri G. The immunopathobiology of syphilis: the manifestations and course of syphilis are determined by the level of delayed-hype hypersensitivity. *Am. J. Dermatopathol*. 2011. Vol.33, No. 5. P. 433-460. DOI: <https://doi.org/10.1097/DAD.0b013e3181e8b587>

19. Dinarello C. Proinflammatory cytokines. *Chest*. 2000. Vol. 118, No. 9. P. 503-508.

DOI: <https://doi.org/10.1378/chest.118.2.503>

20. Kenyon C., Osbak K. K., Crucitti T., Kestens L. The immunological response to syphilis differs by HIV status; a prospective observational cohort study. *BMC Infectious Diseases*. 2017. Vol. 17. P. 111-119. DOI: <https://doi.org/10.1186/s12879-017-2201-7>

21. Knudsen A., Benfield T., Kofoed K. Cytokine expression during syphilis infection in HIV-1-infected individuals. *Sex. Transm. Dis.* 2009. Vol. 36, No. 5. P. 300-304. DOI: <https://doi.org/10.1097/OLQ.0b013e318193ca26>

22. Lusiak M., Podwińska J. Interleukin 10 and its role in the regulation of the cell-mediated immune response in syphilis. *Arch. Immunol Ther. Exp.* 2001. Vol. 49, No. 6. P. 417-421.

Стаття надійшла до редакції
24.06.2019



UDC 618.3-06:[616.24-002.5+616.379-008.64]-07-08

<https://doi.org/10.26641/2307-0404.2019.3.181890>

O.M. Raznatovska¹,
A.V. Fedorec²,
M.O. Shalmina¹,
T.A. Grekova¹

**CLINICAL FEATURES OF PREGNANCY
IN MULTIDRUG-RESISTANT TUBERCULOSIS
AND TYPE 1 DIABETES
MELLITUS COMORBIDITIES
(a case report)**

Zaporizhzhia State Medical University¹

Maiakovskiy av., 26, Zaporizhzhia, 69035, Ukraine

MI «Zaporizhzhia Regional TB Clinical Dispensary»²

Perspektyvna, 2, Zaporizhzhia, 69106, Ukraine

Запорізький державний медичний університет¹

пр. Маяковського, 26, Запоріжжя, 69035, Україна

КУ "Запорізький обласний протитуберкульозний клінічний диспансер" ЗОР²

вул. Перспективна, 2, Запоріжжя, 69106, Україна

e-mail: zsmu@zsmu.zp.ua

Цитування: *Медичні перспективи*. 2019. Т. 24, № 3. С. 101-105

Cited: *Medicni perspektivi*. 2019;24(3):101-105

Key words: pregnancy, multidrug-resistant tuberculosis, type 1 diabetes mellitus

Ключові слова: вагітність, мультирезистентний туберкульоз, цукровий діабет 1-го типу

Ключевые слова: беременность, мультирезистентный туберкулез, сахарный диабет 1-го типа

Abstract. Clinical features of pregnancy in multidrug-resistant tuberculosis and type I diabetes mellitus comorbidities (a case report). Raznatovska O.M., Fedorec A.V., Shalmina M.O., Grekova T.A. *Objective – to update the literature data with the clinical features of pregnancy in multidrug-resistant tuberculosis (MRD-TB) and type 1 diabetes mellitus (T1DM) comorbidities based on an example from own clinical experience. A clinical case of pregnancy course in MRD-TB and T1DM comorbidities was described based on our own clinical experience. We report the clinical case of newly diagnosed MRD-TB in a 38-year-old woman suffering from T1DM. Her general condition was unstable from satisfactory to moderately severe despite an adequate treatment of MRD-TB and T1DM with manifestations of intoxication syndrome and nephropathy. Adenomyosis with periodic bloody vaginal discharge was diagnosed. There was no clinical-radiological dynamics and sputum culture conversion. After an intensive phase of antimycobacterial therapy within 3 months, the patient got pregnant. Based on the medical indications (MRD-TB, absence of sputum culture conversion and clinical-radiological dynamics, moderate severity of T1DM, nephropathy) and adenomyosis with bloody vaginal discharge, the patient was requested to induce the pregnancy termination, and she consented. On month 7 of antimycobacterial therapy, an extensive drug-resistance of Mycobacterium tuberculosis*